

Daftar Pustaka

- [1] McNeil A.J., Frey R., Embrechts P. *Quantitative Risk Management*, 2005.
- [2] Lindskog, F., McNeil, A. and Schmock, U. (2001) *A Note on Kendalls Tau for Elliptical Distribution*. ETH Zurich, working paper
- [3] Kahya, Goksel.B.S (1991). New Modified Anderson-Darling and Cramer-von Mises Goodnessof-fit Tests for a Normal Distribution with Specified Parameters. Ohio.
- [4] Nelsen, R.B. (2006). Second ed. An Introduction to Copulas. Springer, New York.
- [5] Clayton, D.G., 1978. A model for association in bivariate life tables and its application to a uranium exploration data set. *Technometrics* 28, 123-131
- [6] Genest, C., 1987. Franks family of bivariate distribution. *Biometrika* 74, 549-555. Gumbel, E.J., 1960. Bivariate exponential distributions. *Journal of American Statistical Association* 55, 698-707.
- [7] Hanafi, M. 2006. Manajemen Risiko. Yogyakarta: Unit Penerbit Dan Percetakan Sekolah Tinggi Manajemen YKPN.
- [8] Rohmawati,A.A.(2014). Eksplorasi hubungan Value-at-Risk dan Condotional Value-at-Risk. Thesis Magister pada Institut Teknologi Bandung: tidak diterbitkan.

- [9] Danielsson, J. Financial Risk Forecasting The Theory and Practice of Forecasting Market Risk with Implementation in R and Matlab, 2011.
- [10] Artzner, P., F. Delbaen, J. M. Eber dan D. Heath. (1999). Coherent Measures of Risk. Mathematical Finance 9:20322