

REFERENCES

- [1] E. R. Septian and E. N. Hadi, “Reducing Stigma of People with Disabilities: A Systematic Review,” *J. Med. Heal. Stud.*, vol. 2, no. 2, 2021, doi: 10.32996/jmhs.2021.2.2.3.
- [2] H. M. K. K. M. B. Herath and W. R. De Mel, “Controlling an Anatomical Robot Hand Using the Brain-Computer Interface Based on Motor Imagery,” *Adv. Human-Computer Interact.*, vol. 2021, 2021, doi: 10.1155/2021/5515759.
- [3] A. Friant, F. Mulyani, H. Joachim, D. Anggreni, and W. Y. Effendi, “Employment Situation of Person with Disabilities: Case Study in Indonesia,” *Integr. J. Bus. Econ.*, vol. 3, no. 1, 2019, doi: 10.33019/ijbe.v3i1.91.
- [4] N. Birbaumer, “Breaking the silence: Brain-computer interfaces (BCI) for communication and motor control,” in *Psychophysiology*, 2006, vol. 43, no. 6, doi: 10.1111/j.1469-8986.2006.00456.x.
- [5] L. E. H. van Dokkum, T. Ward, and I. Laffont, “Brain computer interfaces for neurorehabilitation-its current status as a rehabilitation strategy post-stroke,” *Ann. Phys. Rehabil. Med.*, vol. 58, no. 1, 2015, doi: 10.1016/j.rehab.2014.09.016.
- [6] N. Robinson, C. Guan, A. P. Vinod, K. Keng Ang, and K. Peng Tee, “Multi-class EEG classification of voluntary hand movement directions,” *J. Neural Eng.*, vol. 10, no. 5, 2013, doi: 10.1088/1741-2560/10/5/056018.
- [7] V. Scarapicchia, C. Brown, C. Mayo, and J. R. Gawryluk, “Functional magnetic resonance imaging and functional near-infrared spectroscopy: Insights from combined recording studies,” *Frontiers in Human Neuroscience*, vol. 11, 2017, doi: 10.3389/fnhum.2017.00419.

- [8] M. Alimardani, S. Nishio, and H. Ishiguro, “Brain-Computer Interface and Motor Imagery Training: The Role of Visual Feedback and Embodiment,” in *Evolving BCI Therapy - Engaging Brain State Dynamics*, 2018.
- [9] M. A. L. Nicolelis *et al.*, “Chronic, multisite, multielectrode recordings in macaque monkeys,” *Proc. Natl. Acad. Sci. U. S. A.*, vol. 100, no. 19, 2003, doi: 10.1073/pnas.1934665100.
- [10] S. Midha, H. A. Maior, M. L. Wilson, and S. Sharples, “Measuring Mental Workload Variations in Office Work Tasks using fNIRS,” *Int. J. Hum. Comput. Stud.*, vol. 147, 2021, doi: 10.1016/j.ijhcs.2020.102580.
- [11] T. Tamura and W. Chen, *Seamless healthcare monitoring: Advancements in wearable, attachable, and invisible devices*. Springer International Publishing, 2017.
- [12] D. Dadebayev, W. W. Goh, and E. X. Tan, “EEG-based emotion recognition: Review of commercial EEG devices and machine learning techniques,” *Journal of King Saud University - Computer and Information Sciences*, vol. 34, no. 7. 2022, doi: 10.1016/j.jksuci.2021.03.009.
- [13] A. Czeszumski *et al.*, “Hyperscanning: A Valid Method to Study Neural Interbrain Underpinnings of Social Interaction,” *Frontiers in Human Neuroscience*, vol. 14. 2020, doi: 10.3389/fnhum.2020.00039.
- [14] H. U. Amin, A. S. Malik, A. R. Subhani, N. Badruddin, and W. T. Chooi, “Dynamics of scalp potential and autonomic nerve activity during intelligence test,” in *Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics)*, 2013, vol. 8226 LNCS, no. PART 1, doi: 10.1007/978-3-642-42054-2_2.
- [15] M. Á. Luján, M. V. Jimeno, J. M. Sotos, J. J. Ricarte, and A. L. Borja, “A survey on eeg signal processing techniques and machine learning: Applications to the neurofeedback of autobiographical memory deficits in schizophrenia,” *Electronics (Switzerland)*, vol. 10, no. 23. 2021, doi: 10.3390/electronics10233037.

- [16] M. Z. Baig, N. Aslam, and H. P. H. Shum, “Filtering techniques for channel selection in motor imagery EEG applications: a survey,” *Artif. Intell. Rev.*, vol. 53, no. 2, 2020, doi: 10.1007/s10462-019-09694-8.
- [17] Abdullah, I. Faye, and M. R. Islam, “EEG Channel Selection Techniques in Motor Imagery Applications: A Review and New Perspectives,” *Bioengineering*, vol. 9, no. 12. 2022, doi: 10.3390/bioengineering9120726.
- [18] I. Elamvazuthi *et al.*, “Development of Electromyography Signal Signature for Forearm Muscle,” in *Procedia Computer Science*, 2015, vol. 76, doi: 10.1016/j.procs.2015.12.347.
- [19] H. Fauzi, M. A. Azzam, M. I. Shapiai, M. Kyoso, U. Khairuddin, and T. Komura, “Energy extraction method for EEG channel selection,” *Telkomnika (Telecommunication Comput. Electron. Control.)*, vol. 17, no. 5, 2019, doi: 10.12928/TELKOMNIKA.v17i5.12805.
- [20] Y. Putranto, “An Improved Performance of Support Vector Machine to Classify EEG Motor Imagery based on Differential Asymmetry,” *PRZEGŁĄD ELEKTROTECHNICZNY*, vol. 1, no. 6, 2023, doi: 10.15199/48.2023.06.40.
- [21] W. Bin Ng, A. Saidatul, Y. F. Chong, and Z. Ibrahim, “PSD-Based Features Extraction for EEG Signal during Typing Task,” in *IOP Conference Series: Materials Science and Engineering*, 2019, vol. 557, no. 1, doi: 10.1088/1757-899X/557/1/012032.
- [22] P. Comon, “Independent component analysis, A new concept?,” *Signal Processing*, vol. 36, no. 3, 1994, doi: 10.1016/0165-1684(94)90029-9.
- [23] A. Tharwat, “Independent component analysis: An introduction,” *Applied Computing and Informatics*, vol. 17, no. 2. 2018, doi: 10.1016/j.aci.2018.08.006.
- [24] I. Marriott Haresign *et al.*, “Automatic classification of ICA components from infant EEG using MARA,” *Dev. Cogn. Neurosci.*, vol. 52, 2021, doi: 10.1016/j.dcn.2021.101024.

- [25] A. H. Khan, M. Hussain, and M. K. Malik, “Arrhythmia Classification Techniques Using Deep Neural Network,” *Complexity*, vol. 2021, 2021, doi: 10.1155/2021/9919588.
- [26] J. S. Rani Alex, M. A. Haque, A. Anand, A. Gautam, and N. Venkatesan, “A deep learning approach for robotic arm control using brain-Computer interface,” *Int. J. Biol. Biomed. Eng.*, vol. 14, 2020, doi: 10.46300/91011.2020.14.18.
- [27] P. Z. Yan, F. Wang, N. Kwok, B. B. Allen, S. Keros, and Z. Grinspan, “Automated spectrographic seizure detection using convolutional neural networks,” *Seizure*, vol. 71, 2019, doi: 10.1016/j.seizure.2019.07.009.